CTA Practical Guide Series, No. 13

For more information:

Market contacts

National Apiculture Development Organisation (TUNADO) P.O. Box 5318 Kampala-Uganda Tel.: +256-41-343160, E-mail: tunadobee@yahoo.co.uk

Honey Care Kenya Limited P.O. Box 24487-00502, Nairobi, Kenya. Tel/Fax: +254-20-3874448/50 Email: info@honeycareafrica.com, Website: www.honeycareafrica.com

Africa Now Kenya Limited P.O. Box 2514, Kisumu Kenya Tel/Fax: +254-57-2021181 E-mail: kisumu@africanow.org, Website: www.africanow.org

Training contacts

Bee Natural Products Uganda Ltd P.O. Box 5318, Pan Africa House, Plot 3 Kimathi Avenue Kampala, Uganda Tel: +256-41-253844 or 234676; Fax: +256-41-259052 Website: www.beenaturalproducts.com

International Centre for Insect Physiology and Ecology (ICIPE) P.O. Box 30772-00100, Nairobi, Kenya Tel: +254 (20) 8632000; Fax: +254 (20) 8632001/8632002 E-mail: icipe@icipe.org; Website: www.icipe.org

National Agricultural Advisory Services (NAADS) P.O. Box 25235, Kampala, Uganda Tel: +256-41-345440/345065/345066; Fax: +256-41-347843 E-mail: naads@iwayafrica.com; Website: www.naads.or.ug

> Sasakawa Global 2000 (MoA) P.O. Box 62347,Addis Ababa Ethiopia Tel: +251-115-518040; Fax: +251-115-512984 E-mail: moa@telecom.net.et



partageons les connaissances au profit des communautés rurales sharing knowledge, improving rural livelihoods

The ACP-EU Technical Centre for Agricultural and Rural Cooperation (CTA) P.O. Box 380, 6700 AJ Wageningen, The Netherlands, Tel: +31(0)317-467100, Fax: +31(0)317-460067, E-mail: cta@cta.int, Website: www.cta.int

CTA is financed by the European Union



© CTA 2007 - ISSN 1873-8192 The information in this guide can be freely reproduced for non-commercial use, if credited as coming from CTA. Reproduction for commercial use requires prior authorization from CTA

How to Keep Bees and Process Honey



How to Keep Bees and Process Honey


Asaf Ayatuhaire, from Bushenyi District in South Western Uganda, has more than 100 beehives of different types.With an annual production of honey of around 2,000 kilogrammes, he earns more than US\$ 2,200 a year and has recently registered a company,West Honeys Uganda.

But ten years ago, Asaf had just three traditional hives and produced only six kilogrammes of honey a year. Then he attended a national training of trainers programme, organized and sponsored by the Entomology Department of the Ministry of Agriculture. After completing his training, he became involved in training other beekeepers in his local communities. Asaf was also able to interact with NGOs and international and local funding agencies, such as USAID, FAO and the District Farmers Association. These agencies helped him to expand his apiary and acquire improved top-bar hives.

Today, thanks to beekeeping, Asaf has a permanent house, pays fees for his children in secondary and higher education and is able to save from his income.

Why keep bees?

- Bees pollinate crops and thus help increase yields
- High demand for honey exists in local, regional and international markets
- Propolis, collected from plants by bees to cover the inside of the hive, treats a broad range of ailments
- Pollen, collected from plants by bees to feed their larvae, is used in the perfume industry, and is a food additive and a medicine
- Royal jelly, made by young bees from gland secretions and fed to the queen to make her strong, has medicinal properties



Freshly harvested honey comb



- · Beewax is used in cosmetics, candles and polishes
- Beekeeping has low start-up costs and occupies negligible land space
- About 80% of honey consists of sugars that are readily absorbed by the body and honey is thus quite suitable for children, sick people and those who perform heavy manual tasks.

Choosing the right hive

The table below lists the advantages and disadvantages of three different types of hive.

	Type of hive	Advantages	Disadvantages
	Local woven beehive	Cheap and easy to establish Made from locally available materials	The colony cannot be inspected
			It is difficult to prevent swarming
			or replace a queen
			Quality cannot be controlled because the combs with brood
		More propolis is produced than in other types of hive	cannot be separated from the
			honeycombs
			Honey yields are low
			(approximately 6 to 10 kg per year)
			Life span of the hive is up to 2
	t		years

Type of hive	Advantages	Disadvantages
Top-bar hive	Can be made with locally available and inexpensive materials	More expensive than local hives
		Difficult for small-scale farmers to acquire minimum economic
	Easy inspection and control of swarming	number (about 10 hives)
		Predators and other insects can easily access them because they are hung low
	Quality of honey can be monitored	
	Yields are 20 to 40 kg per year	
	Life span is about 10 years	
	More wax is produced since the combs have to be cut from the frame during harvesting	
Langstroth hive	Combs with brood (young bees) can be easily separated from combs with honey	Costly to build or buy
		Requires a special extractor to remove the honey from the combs
	Yields range between 50 and 60 kg per year	Needs more advanced beekeeping skills
	Quality of honey can be monitored	

Site selection

Beehives should be placed near a source of water (river, stream, lake, dam or natural or man-made ponds) and in an area with adequate vegetation, such as coffee, sunflower, moringa, mangoes, oranges, bananas and other flowering plants.

The site should be fenced to protect bees from people and animals (and people and animals from bees).

Hives should be located at least 30 metres away from roads and public or noisy places.

Populating the hive

There are two ways of populating a swarm:

I. Baiting a swarm

- Use a small hive that has already been inhabited by bees and fill it with frames or top bars; two of the frames should contain combs and the others should have foundation sheets or strips of old comb.
- Place the hive in a tree or on a roof in such a way that there is some protection from the wind.
- Place the hive in its desired place the very day that the swarm has taken occupation of it.
- Baiting is useful only in the swarming season, usually at the beginning of the dry season and end of a cold season.

2. Capturing a swarm

You may capture a newly settled swarm around a branch of a tree, but make sure you wear protective clothing.


- First sprinkle cold water over the bees with a brush or spray bottle to stop them from moving away.
- Hold a small hive without frames, a basket or a box under the swarm.
- Drive the bees into the hive with a bee brush or smoke.

Honey harvesting

For traditional and top-bar hives, harvest once or twice a year. For the Langstroth hive, harvest three or four times a year.

Steps in harvesting honey:

- Wear protective clothing: overall, veils, gloves, boots
- Light smoker
- Gently open the hive
- Select combs that are two-thirds sealed full of honey
- Avoid combs with brood
- Lift comb, blow smoke on both sides and gently brush bees back into hive with a feather
- Cut away combs and honey, leaving about 1 cm of comb on the bar
- Place capped honey in a clean, dry container and cover
- Leave at least eight combs for the bees
- Before closing hive, push the unripe combs next to the combs with brood and place harvested bars behind these.






2 harvesting process

Processing of honey

- Use a hot rod or knife to de-cap the honeycombs (by passing it over the combs).
- Place the de-capped comb on a piece of fine white linen tied over the top of a plastic container. The de-capped side should face downwards.
- Let the honey drip through the cloth until the cells are empty.
- Turn over the comb and repeat the above process.
- Pack honey in clean, dry jars with well-sealed lids. Label with date harvested and store in a cool place.



3 harvested honey in comb

market.

It is important to keep records during each hive inspection in order to follow the progress of each colony and monitor its condition. Hives should be inspected two to three-times a month. Entries may be made under the following headings: hive number, date hive occupied, date of inspection, observation, yield in kilogrammes.

What can go wrong	Cause	Solution
Bees fly away	Lack of food and water Disturbance of hive e.g. by fire, people or animals Frequent attacks by pests	Ensure sufficient food: plant flowering plants near hive Provide water during droughts Reduce size of colony Place hive at least 30 metres from roads and public places and fence site Treat bees gently Keep hive area clean
Bee sting	Disturbance of hive	Keep calm, remove the sting and smoke the stung area In case of severe reaction, report to nearest hospital or clinic
Pests (safari ants, wax moth, honey badger)	Poor siting and maintenance of hives	Hang hives between trees If poles are used, treat poles with used engine oil or grease to keep out termites and ants
Use of insecticides or pesticides which poison bees	Use of chemical sprays nearby	Keep in touch with other farmers and know when they are spraying Request farmers to spray at night when bees are safely in hive
Fires	Bush burning	Clear grass and bushes surrounding the apiary